


26 June 1993 No1879 Weekly £1.50

ISSN 0262 4079

NewScientist



**The trouble
with implants**

**Last gasp of the
Namib baboons**

**Science v. Art...
the story continues**

Reviving the Rhine



NewScientist

Vol 138 No 1879

26 June 1993

TECHNION
ISRAELI INSTITUTE OF TECHNOLOGY
ESTABLISHED IN 1956
FINANCED BY
THE RASHI FOUNDATION

NEWS

THIS WEEK 4

Onco-mouse flops/Mapping Muttley/Making war with information/Hubble rescue/Britain axes Antarctic bases/Shetland escapes disaster/Europe agrees forest principles/Alaska hunts down its wolves/In Brief

FOCUS Carbon dioxide's taxing questions 12

Fred Pearce
Can Britain and the other industrial nations meet the deadline agreed a year ago for stabilising emissions of carbon dioxide? And do they need to do more?

SCIENCE 14

Edge of the Solar System/Milky Way's dark halo/Introns: DNA's error-checkers/Reading ant trails/Cambrian motorbikes

TECHNOLOGY 18

Paperless phone books/Star Wars on graffiti/Easier timetabling/Recycling tyres/Hybrid cars/Patents

VIEWPOINTS

COMMENT 3

REVIEW 39

Revisionists welcome, ranters not/Africa's fearsome future/Apes' rights/Genetics illuminated/A matter of life and death/Moon rocks/24-hour days/Right writing

FORUM 44

Who says they can forecast the weather?/Can do, therefore must do

Thistle Diary 46

Enigma

Life, the Universe 47

and (almost) everything

LETTERS 48

FEEDBACK 80

APPOINTMENTS 50




Amy Guip

Shaping up with silicone: are breast implants making women ill? p 22



Bruce Coleman

Waiting for water: survival in the desert, p 34



In rude health: but what if the old have had enough of living? p 41

ZIFA



Ferrero/Labat/Ardea

Hunter hunted: Alaska's plan to kill wolves, p 9

FEATURES

COVER The trouble with implants 22

Susan Katz Miller
For thirty years plastic surgeons said silicone implants were safe. Now women are rushing to have them removed as researchers explore the links with disease

Greenprint for rescuing the Rhine 25

Fred Pearce
Orthodox methods for cleaning up the Rhine have only been partially successful. Now the nations responsible are starting to back the ecologists' vision of a river that is fit for salmon

A tale of two cultures 30

Clive Rassam
In 1959 C. P. Snow coined a phrase "the two cultures" to describe what he saw as a widening gap between science and the arts. The debate still rumbles on, but are Snow's two cultures more perceived than real?

Real monkeys don't drink water 34

Sue Armstrong
A remarkable troop of southern African baboons has learnt how to survive in a desert unfit for primate life. But their days could be numbered

This week's cover is by Amy Guip

NEXT WEEK

Did Tibet cool the world?
The collision of Asia and India more than 40 million years ago could be responsible for today's climate

How to stop head-banging
Materials made to measure NASA's shrinking space station

Published by
ipcmagazines
Distributed by IPC Marketforce

New Scientist is published weekly by IPC Magazines Ltd, part of Reed Publishing Europe. ISSN No 0262 4079. Registered at the Post Office as a newspaper and printed in England. An annual subscription in the USA is US \$140 including airspeed delivery. 2nd class postage paid at Newark NJ, USA POSTMASTER: Send address changes to New Scientist, c/o Virgin Mailing & Distribution, Cargo Building 150, Newark International Airport, New Jersey, NJ 07114. Airfreight and mailing in the USA by Publications Expediting Inc, 200 Meacham Avenue, Elmont, NY 11003. © 1993 IPC Magazines Limited, England.

Mathematicians learn how to read the ant-trails

Ian Stewart

"GO TO the ant, thou sluggard; consider her ways, and be wise," says *Proverbs*, vi, 6. Although the writer of this line was probably thinking of ants' legendary dedication to hard work, the ways of ants have a lot more to teach the seeker after wisdom—for instance, how to find the shortest route between two points without having a birds' eye view of the terrain. Now a mathematician in Israel, studying "pursuit problems", has discovered how ants can achieve the feat by repeatedly applying simple local rules.

As a "scout" ant searches for food, it leaves a faint trail of pheromones. Other ants that follow the trail lay down their own pheromones, making the trail stronger and straighter. Because individuals, such as the scout itself, move rather randomly, it is hard to see how the ants collectively "know" the shortest path.


Many people, including the physicist Richard Feynman, have wondered at their ability. The consensus is that the straight path ants settle on is a result of a sequence of successive improvements. Although the route taken by the scout is random, the ants that follow modify the route and make it more efficient. There is evidence that ants that are forced to make sharp changes in direction either perform a U-turn, or keep going but lay down less pheromone. Computer simulations show that these simple rules for ant behaviour lead to a build-up of pheromone along tracks that are relatively straight (*New Scientist*, Science, 27 March).

Feynman's explanation was similar. He thought that ants trying to follow the original, random path tend to overshoot at corners and wander off randomly until they meet the path again. The net effect is to eliminate sharp bends and to cause pheromone to accumulate along straight-line paths. This explanation is reminiscent of the quantum explanation of why—in classical physics—light always takes the shortest path between any two points. In the quantum world, the light follows all possible paths, but when these are superposed, the wave-functions interfere destructively away from the classical path. Feynman pioneered this kind of explanation in physics, so it is not surprising that he developed a similar theory for ant trails: randomly moving ants superpose their pheromones, and the peak concentration is along the straight-line path.

Physical analogies and computer experiments are all very well, but mathematicians prefer proofs. Alfred Bruckstein of Technion University in Haifa, Israel, set up a rigorous mathematical model that implements a



Kim Taylor/Bruce Coleman



Robert Paz/Caltech

Richard Feynman: saw the deep connections between ants and photons, between pheromones and wave functions

particular version of Feynman's suggestion. He proved that as successive ants follow each other, their paths become ever more straight (*The Mathematical Intelligencer*, Spring 1993, p 59).

Bruckstein's model replaces the pheromones by dynamical rules, but this does not deny the importance of pheromones in practice: the rules are a simplified version of "follow the previous pheromone trail but smooth it out a bit".

The model lies in the general area of "pursuit problems", which date back to Leonardo da Vinci. A number of objects—

usually represented by mathematical points—chase each other, so that at each instant each object is moving directly towards some other chosen object. For example, four beetles start from the corners of a square, and each one chases the next. They spiral in and meet at the centre in the same time that it would take one beetle to traverse the edge of the square. A basic pursuit theorem states that if several objects chase each other in sequence (so that A chases B chases C chases D chases A, say) then eventually they will all meet at the same point.

In Bruckstein's model, one scout ant sets off and lays down a trail. A second ant, moving at the same speed, starts some distance behind, but obeys the pursuit rule: "Always head towards the scout." A third ant pursues the second in the same manner, and so on. The motion of each ant is determined by a differential equation, which states that its velocity vector is always directed towards the current position of the previous ant.

Bruckstein considered the angle through which a given ant had turned at a given time, and estimated how this angle changes from one trip to the next. He found that the angle tended to zero, implying that the paths become ever more straight.

Bruckstein's theorem should be seen as a contribution to the mathematics of pursuit, motivated by ant trails, rather than a theory of ant behaviour. Real ants wander at varying speeds along a pheromone-splashed landscape, rather than playing follow-the-leader at uniform speeds. Indeed, it would be interesting to model the deposition of pheromones more directly: the analogy with quantum mechanics is quite strong and techniques similar to Feynman path integrals—which superpose contributions from all possible paths—might be useful.

Be that as it may, Bruckstein's theorem provides an elegant example of how systems that obey purely local rules can collectively settle into global optima. Many cases of collective animal behaviour, such as the flocking of birds or the movements of shoals of fish, display surprising large-scale patterns, giving the impression that the entire flock or shoal has some kind of overall "consciousness" or sense of purpose. In fact, this appearance is an illusion—it is a collective response arising from the combination of simple rules followed by each individual. The important thing about the ant-trail pursuit problem is that for once we can pin down mathematically the precise manner in which the collective pattern is generated. □